

Natural variability of restriction profiles in non-coding part of *Prunus persica* (L.) Batsch. Pru p 3 gene

Lucia Zamiešková*, Jana Žiarovská, Jana Bilčíková, Veronika Fialková
Slovak University of Agriculture in Nitra, Slovakia

References

- Ahrazem, O. et al. (2007). Assessing allergen levels in peach and nectarine cultivars. *Annals of Allergy Asthma and Immunology*, 99(1), 42-47.
- Alshamary, S.F. and Al-Horayess, O.S. (2013). Appraisal of mineral and heavy metal contents in peach and grape grown at some major agricultural companies in Saudi Arabia. *Oriental Journal of Chemistry*, 29(4), 1515.
- Aranzana, M.J. et al. (2019). Prunus genetics and applications after de novo genome sequencing: achievements and prospects. *Horticulture research*, 6 (58). <https://doi.org/10.1038/s41438-019-0140-8>
- Aranzana, M.J. et al. (2010). Genetic variation, population structure and linkage disequilibrium in peach commercial varieties. *BMC Genetics*, 11(1), 69.
- Baird, W.V. et al. (1996). Progress in Prunus mapping and application of molecular markers to germplasm improvement. *Horticulture Science*, 31(7), 748-749.
- Bolhaar, S. et al. (2005). In vivo assessment with prick-to-prick testing and double-blind, placebo controlled food challenge of allergenicity of apple cultivars. *Journal of Allergy and Clinical Immunology*, 116(5), 1080-1086.
- Borges, J.P. et al. (2006). The lipid transfer proteins (LTP) essentially concentrate in the skin of Rosaceae fruits as cell surface exposed allergens. *Plant Physiology and Biochemistry*, 44(10), 535-542.
- Botton A., et al. (2002) Differential expression of two lipid transfer protein genes in reproductive organs of peach (*Prunus persica* L. Batsch). *Plant Science*, 163, 993-1000.
- Botton, A. et al. (2009). Peach (*Prunus persica* L. Batsch) allergen-encoding genes are developmentally regulated and affected by fruit load and light radiation. *Journal of Agricultural and Food Chemistry*, 57(2), 724-734.
- Bouhadida, M. et al. (2010). Genetic variability of introduced and local Spanish peach cultivars determined by SSR markers. *Tree Genetics and Genomes*, 7(2), 257-270.
- Brenna, O.V. et al. (2004). Presence of Allergenic Proteins in Different Peach (*Prunus persica*) Cultivars and Dependence of Their Content on Fruit Ripening. *Journal of Agriculture and Food Chemistry*, 52(26), 7997-8000.
- Carnés, J. et al. (2002). Pru p 3 (LTP) content in peach extracts. *Allergy*, 57(11), 1071-1075.
- Chae, K. et al. (2009) . A gain of function mutation of Arabidopsis lipid transfer protein 5 disturbs pollen tube tip growth and fertilization. *Plant Cell*, 21(12), 3902-3914.
- Chen, L., et al. (2008) Genomic characterization of putative allergen genes in peach/almond and their synteny with apple. *BMC Genomics*, 9 (1), 543.
- Cheng, Z. and Huang, H. (2009) . SSR fingerprinting Chinese peach cultivars and landraces (*Prunus persica*) and analysis of their genetic relationships. *Science Horticulture*, 120(2), 188-193.
- Cubells-Baeza, N. et al. (2017). Identification of the ligand of Pru p 3, a peach LTP. *Plant Molecular Biology*, 94(1-2), 33-44.
- Duffort, O.A. et al. (2002) Immunoassay to quantify the major peach allergen Pru p 3 in foodstuffs. Differential allergen release and stability under physiological conditions. *Journal of Agriculture and Food Chemistry*, 50(26), 7738-7741.
- Fernández-Rivas, M. et al. (2003). Clinically relevant peach allergy is related to peach lipid transfer protein, Pru p 3, in the Spanish population. *Journal of Allergy and Clinical Immunology*, 112(4), 789-795.
- Fernández-Rivas, M. et al. (2006). Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. *Journal of Allergy and Clinical Immunology*, 118(2), 481-488.

* **Corresponding Author:** Lucia Zamiešková, Slovak University of Agriculture in Nitra, Department of Genetics and Plant Breeding, Tr. A. Hlinku 2, 94976 Nitra, Slovak Republic. E-mail: z zamieskova@uniag.sk

- Gamboa, P.M. et al. (2007). Two different profiles of peach allergy in the north of Spain. *Allergy*, 62(4), 408-414.
- Gao, Z.S. et al. (2008). Assessment of allelic diversity in intron-containing Mal d 1 genes and their association to apple allergenicity. *BMC Plant Biology*, 13(8), 116.
- García-Casado, G. et al. (2003). Identification of IgE-binding epitopes of the major peach allergen Pru p 3. *Journal of Allergy and Clinical Immunology*, 112(3), 599-605.
- Helsper, J. et al. (2002). Quadrupole time-of-flight mass spectrometry: a method to study the actual expression of allergen isoforms identified by PCR cloning. *Journal of Allergy and Clinical Immunology*, 110(1), 131–138.
- Hoffmann-Sommergruber, K. and SAFE consortium. (2005). The SAFE project: 'plant food allergies: field to table strategies for reducing their incidence in Europe' an EC-funded study. *Allergy*, 60(4), 436-442.
- Mitic, S. et al. (2019). Comparative Study on Minerals in Peel and Pulp of Peach (*Prunus persica* L.) Fruit. *Revista de Chimie Bucharest Original Edition*, 7(6), 2281-2285.
- Monaci, L. and Visconti, A. (2009). Mass spectrometry-based proteomics methods for analysis of food allergens Tr. In TrAC Trends. *Analytical Chemistry*, 28(5), 581-591.
- Pasquato, N. et al. (2006). Crystal structure of peach Pru p 3, the prototypic member of the family of plant non-specific lipid transfer protein panallergens. *Journal of Molecular Biology*, 356(3), 684-694.
- Pastorello, E.A. et al. (1999). The major allergen of peach (*Prunus persica*) is a lipid transfer protein. *Journal of Allergy and Clinical Immunology*, 103(3), 520-526.
- Reuter, A. et al. (2006). A critical assessment of allergen component-based *in vitro* diagnosis in cherry allergy across Europe. *Clinical and Experimental Allergy*, 36(6), 815-823.
- Salcedo, G. et al. (2007). Plant non-specific lipid transfer proteins: an interface between plant defence and human allergy. *Biochimistry Biophysical Acta*, 1771(6), 781-791.
- Salcedo, G. et al. (2008). Review. Plant food allergens: peach non-specific lipid transfer protein Pru p 3 as a model. *Spanish Journal of Agricultural Research*, 6(Special issue), 30-37.
- Sancho, A.I. et al. (2008). Measurement of lipid transfer protein in 88 apple cultivars. *International Archives of Allergy and Immunology*, 146(1), 19–26.
- Verde, I. et al. (2013). The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nature Genetics*, 45pr(5), 487- 494.
- Wills, R. B. H. et al. (1983). Nutrient composition of stone fruit (*Prunus* spp.) cultivars: Apricot, cherry, nectarine, peach and plum. *Journal of the Science of Food and Agriculture*, 34(12), 1383–1389.
- Wünsch, A. et al. (2005). Molecular characterization of local Spanish peach [*Prunus persica* (L.) Batsch] germplasm. *Genetic Resources and Crop Evolution*, 53(5), 925–932.
- Xie, R. et al. (2010). Evaluation of the genetic diversity of Asian peach accessions using a selected set of SSR markers. *Science Horticulture*, 125(4), 622–629.
- Yang, Z., et al. (2011). Differential transcript abundance and genotypic variation of four putative allergen-encoding gene families in melting peach. *Tree Genetics & Genomes*, 7(5), 903.
- Ying-Tao, M. et al. (2014). The PpLTP1 Primary Allergen Gene is Highly Conserved in Peach and Has Small Variations in Other *Prunus* Species. *Plant Molecular Biology Reporter*, 32(3), 652-663.
- Zhang, A.M. et al. (2006). Genetic diversity and ecogeographical phylogenetic relationships among peach and nectarine cultivars based on simple sequence repeat (SSR) markers. *Journal of American Society of Horticulture Science*, 131(4), 513–521.
- Zhong-Shan, G. et al. (2016). Quantification of Peach Fruit Allergen Lipid Transfer Protein by a Double Monoclonal Antibody-based Sandwich ELISA. *Food Analytical Methods*, 9(4), 823–830.
- Zuidmeer, L. and Van Ree, R. (2007). Lipid transfer protein allergy: primary food allergy or pollen/food syndrome in same cases. *Current Opinion of Allergy and Clinical Immunology*, 7(3), 269-273.