

The effect of essential oils on quality and mineral composition of shelleggs

Róbert Herkeľ, Branislav Gálik, Daniel Bíro, Michal Rolinec, Milan Šimko, Miroslav Juráček, Henrieta Arpášová, Ondrej Hanušovský

References

- ANDJELKOVIC, M. et al. (2010) Phenolic compounds and some quality parameters of pumpkin seed oil. *European Journal of Lipid Science and Technology*, vol. 112, pp. 208-217. doi:<http://dx.doi.org/10.1002/ejlt.200900021>
- AOAC. (2000). *Official methods of analysis*. Washington: Association of official analytical chemists.
- ARPÁŠOVÁ, H. et al. (2014) The influence of oregano essential oil and *Rhus coriaria* L. on qualitative parameters and microbiological indicators of hens eggs content. *Scientific Papers: Animal Science and Biotechnologies*, vol. 47, no. 2, pp. 6-12.
- ARPÁŠOVÁ, H. et al. (2015) The effect of selected feed additives on the shell qualitative parameters of table eggs. *Scientific Papers: Animal Science and Biotechnologies*, vol. 48, no. 1, pp. 6-10.
- AYDIN, R., PARIZA, M. W. and COOK, M. E. (2001) Olive oil prevents the adverse effects of dietary conjugated linoleic acid on chick hatchability and egg quality. *The Journal of Nutrition*, vol. 131, no. 3, pp. 800-806.
- BAR, A., VAX, E. and STRIEM, S. (1999) Relationships among age eggshell thickness and vitamin D metabolism and its expression in the laying hen. *Comparative Biochemistry and Physiology – Part A: Molecular and Integrative Physiology*, vol. 123, pp. 147-154.
- BOTSOGLOU, N. et al. (2005) The effect of feeding rosemary, oregano, saffron and alpha-tocopheryl acetate on hen performance and oxidative stability of eggs. *South African Journal of Animal Science*, vol. 35, no. 3, pp. 143-151.
- European Commission (2007): Council Regulation (EC) No 834/2007 of 28 June 2007 on organic production and labelling of organic and repealing regulation (EEC), 2092/91. *Official Journal*, L 189, 20/07/2007, pp.1-23.
- FAITARONE, A. B. G. et al. (2013) Cholesterol levels and nutritional composition of commercial layers eggs fed diets with different vegetable oils. *Brazilian Journal of Poultry Science*, vol. 15, no. 1, pp. 31-38.
- HASHEMI, J. M. (2013) Pumpkin seed oil and vitamin E improve reproductive function of male rats inflicted by testicular injury. *World Applied Sciences Journal*, vol. 23, no. 10, pp. 1351-1359. doi:<http://dx.doi.org/10.5829/idosi.wasj.2013.23.10.13153>
- HERKEL', R. et al. (2016) The effect of a phyto-genic additive on nutritional composition of turkey meat. *Journal of Central European Agriculture*, vol. 17, no. 1, pp. 25-39. doi:<http://dx.doi.org/10.5513/JCEA01/17.1.1664>
- CHERIAN, G., CAMPBELL, A. and PARKER, T. (2009) Egg quality and lipid composition of eggs from hens fed *Camelina sativa*. *Journal of Applied Poultry Research*, vol. 18, pp.143-150. doi:<http://dx.doi.org/10.3382/japr.2008-00070>
- CHETTY, K. N. et al. (2004) Garlic induced alteration in liver mineral concentrations in corn oil and olive oil fed rats. *Pathophysiology*, vol. 11, pp. 129-131.
- KÓŇA, J., ĎUROVKA, M. and TANCÍK, J. (2007) *Pumpkin vegetables*. Nitra: Garmond. 148 p.
- KUCUKERSAN, K., YESILBAG, D. and KUCUKERSAN, S. (2010) Influence of different dietary oil sources on performance and cholesterol content of egg yolk in laying hens. *Journal of Biological and Environmental Sciences*, vol. 4, pp. 117-122.
- LI-CHAN, E. C. Y. and KIM, H. O. (2008) Structure and chemical composition of eggs. In: MINE, Y (Ed.). *Egg bioscience and biotechnology*, Wiley-Interscience, Hoboken, pp. 1-8.

- LOKAEWMANEE, K. et al. (2014) Eggshell quality, eggshell structure and small intestinal histology in laying hens fed dietary Pantoea-6 and plant extracts. *Italian Journal of Animal Science*, vol. 13, pp. 332-339. doi:<http://dx.doi.org/10.4081/ijas.2014.3163>
- NAGY, J. et al. (2009) *Hygiene of poultry meat, eggs and venison*. Košice: Editorial Centre of University of Veterinary Medicine, pp. 291–338.
- NYS, Y. et al. (1999) Avian eggshell mineralization. *Poultry and Avian Biology Reviews*, vol. 10, pp. 143-166.
- PANDA, A., RAMA, R. S. and RAJU, M. (2009) Phytobiotics, a natural growth promoter. *Poultry international*, vol. 48, no. 7, pp. 10-11.
- PARK, J. H., UPADHAYA, S. D. and KIM, I. H. (2015) Effect of dietary Marine Microalgae (*Schizochytrium*) powder on egg production, blood lipid profiles, egg quality, and fatty acid composition of egg yolk in layers. *Asian Australasian Journal of Animal Science*, vol. 28, no. 3, pp. 391-397. doi:<http://dx.doi.org/10.5713/ajas.14.0463>
- SHAKOOR, H. I. et al. (2002) Effect of feeding canola and soybean oils on serum lipid profile in commercial layers. *Pakistan Veterinary Journal*, vol. 22, pp. 48-51.
- SHAKOOR, H. I. et al. (2003) Comparative study on the effects of feeding canola and soybean oils on egg production and cholesterol in commercial layers. *Pakistan Veterinary Journal*, vol. 23, pp. 22-26.
- STEVENSON, D. G. et al. (2007) Oil and tocopherol content and composition of pumpkin seed oil in 12 cultivars. *Journal of Agricultural and Food Chemistry*. vol. 55, pp. 4005-4013.
- SUGINO, H., NITODA, T. and JUNEJA, L. R. (1997) General chemical composition of hen eggs. In: YAMAMOTO, T. et al. (eds.). *Hen eggs: Their basic and applied science*. New York: CRC Press, pp. 13-24.
- TŮMOVÁ, E., GOUS, R. M. and TYLER, N. (2014) Effect of hen age, environmental temperature, and oviposition time on egg shell quality and egg shell and serum mineral contents in laying and broiler breeder hens. *Czech Journal of Animal Science*, vol. 59, no. 9, pp. 435-443.
- ZELENKA, J. et al. (2008) The effect of dietary linseed oils with different fatty acid pattern on the content of fatty acids in chicken meat. *Veterinarni Medicina*, no. 2, pp. 77-85.
- ZHANG, Z. F. and KIM, I. H. (2014) Effects of dietary olive oil on egg quality, serum cholesterol characteristics, and yolk fatty acid concentrations in laying hens. *Journal of Applied Animal Research*, vol. 42, pp. 233-237. doi:<http://dx.doi.org/10.1080/09712119.2013.822815>
- ZHAO, G. et al. (2007) Dietary α -linolenic acid inhibits proinflammatory cytokine production by peripheral blood mononuclear cells in hypercholesterolemic subjects. *The American Journal of Clinical Nutrition*, vol. 85, no. 2, pp. 385–391.