

## Assessing footprints of natural selection through PCA analysis in cattle

Nina Moravčíková, Veronika Kukučková, Gábor Mészáros, Johann Sölkner, Ondrej Kadlečík, Radovan Kasarda

### References

AKEY, J. M. et al. (2002) Interrogating a high-density SNP map for signatures of natural selection. *Genome Research*, vol. 12, pp. 1805-1814. doi:<http://dx.doi.org/10.1101/gr.631202>

ALLENDORF, F. W., HOHENLOHE, P. A. and LUIKART, G. (2010) Genomics and the future of conservation genetics. *Nature Reviews Genetics*, vol. 11, no. 10, pp. 697-709. doi:<http://dx.doi.org/10.1038/nrg2844>

BIERNE, N., ROZE, D. and WELCH, J. J. (2013) Pervasive selection or is it . . .? Why are  $F_{ST}$  outliers sometimes so frequent?. *Molecular Ecology*, vol. 22, pp. 2061-2064. doi:<http://dx.doi.org/10.1111/mec.12241>

DRUET, T. et al. (2013). Identification of large selective sweeps associated with major genes in cattle. *Animal Genetics*, vol. 44, pp. 758-762. doi:<http://dx.doi.org/10.1111/age.12073>

DUFORET-FREBOURG, N. et al. (2015) Detecting genomic signatures of natural selection with principal component analysis: application to the 1000 genomes data. *Molecular biology and evolution*, vol. 33, pp. 1082-1093. doi:<http://dx.doi.org/10.1093/molbev/msv334>

DUFORET-FREBOURG, N., BAZIN, E. and BLUM, M. G. B. (2014) Genome scans for detecting footprints of local adaptation using a bayesian factor model. *Molecular biology and evolution*, vol. 31, pp. 2483-2495. doi:<http://dx.doi.org/10.1093/molbev/msu182>

FERENČAKOVIĆ, M., SOLKNER, J. and CURIK, I. (2013) Estimating autozygosity from high-throughput information: effects of SNP density and genotyping errors. *Genetic Selection Evolution*, vol. 45, no. 1, pp. 42. doi:<http://dx.doi.org/10.1186/1297-9686-45-42>

Flori, L. et al. (2009) The genome response to artificial selection: a case study in dairy cattle. *PLoS One*, vol. 4, e6595. doi:<http://dx.doi.org/10.1371/journal.pone.0006595>

GIUSTI, J. et al. (2013) Expression of genes related to quality of *Longissimus dorsi* muscle meat in Nellore (*Bos indicus*) and Canchim (5/8 *Bos taurus* × 3/8 *Bos indicus*) cattle. *Meat Science*, vol. 94, no. 2, pp. 247-252. doi:<http://dx.doi.org/10.1016/j.meatsci.2013.02.006>

GOWANE, G. R. et al. (2014) The Expression of IL6 and 21 in Crossbred Calves Upregulated by Inactivated Trivalent FMD Vaccine. *Animal Biotechnology*, vol. 25, no. 2, pp. 108-118. doi:<http://dx.doi.org/10.1080/10495398.2013.834826>

GUTIÉRREZ-GIL, B., ARRANZ, J. J. and WIENER, P. (2015) An interpretive review of selective sweep studies in *Bos taurus* cattle populations: identification of unique and shared selection signals across breeds. *Frontiers genetics*, vol. 6, pp. 167. doi:<http://dx.doi.org/10.3389/fgene.2015.00167>

LUU, K., BAZIN, E. and BLUM, M. G. B. (2016) pcadapt: An R package for performing genome scans for selection based on principal component analysis. *bioRxiv*. doi:<http://dx.doi.org/10.1101/056135>

MANCINI, G. et al. (2014) Signatures of selection in five Italian cattle breeds detected by a 54K SNP panel. *Molecular Biology Reports*, vol. 41, pp. 957-965. doi:<http://dx.doi.org/10.1007/s11033-013-2940-5>

MARTINS, H. et al. (2016) Identifying outlier loci in admixed and in continuous populations using ancestral population differentiation statistics. *bioRxiv*, p. 054585. doi:<http://dx.doi.org/10.1101/054585>

MCCLURE, M. C. et al. (2012) Genome-wide association analysis for quantitative trait loci influencing Warner–Bratzler shear force in five taurine cattle breeds. *Animal Genetics*, vol. 43, no. 6, pp. 662-673. doi:<http://dx.doi.org/10.1111/j.1365-2052.2012.02323.x>

NIELSEN, R. (2005) Molecular signatures of natural selection. *Annual Review of Genetics*, vol. 39, pp. 197-218. doi:<http://dx.doi.org/10.1146/annurev.genet.39.073003.112420>

NOVEMBRE, J. et al. (2008) Genes mirror geography within Europe. *Nature*, vol. 456, pp. 98-101. doi:<http://dx.doi.org/10.1038/nature07566>

OLEKSYK, T. K., SMITH, M. W. and O'BRIEN, S. J. (2010) Genome-wide scan for footprints of natural selection. *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 365, pp. 185-205. doi:<http://dx.doi.org/10.1098/rstb.2009.0219>

QANBARI, S. et al. (2011) Application of site and haplotype-frequency based approaches for detecting selection signatures in cattle. *BMC Genomics*, vol. 12, pp. 318. doi:<http://dx.doi.org/10.1186/1471-2164-12-318>

STELLA, A. et al. (2010) Identification of selection signatures in cattle breeds selected for dairy production. *Genetics*, vol. 185, pp. 1451-1461. doi:<http://dx.doi.org/10.1534/genetics.110.116111>

STOREY, J. D. (2002) A direct approach to false discovery rates. *Journal of the Royal Statistical Society, Series B*, vol. 64, pp. 479-498. doi:<http://dx.doi.org/10.1111/1467-9868.00346>

WAPLES, R. S. and GAGGIOTTI, O. (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, vol. 15, pp. 1419-1439. doi:<http://dx.doi.org/10.1111/j.1365-294x.2006.02890.x>

WEIR, B. S. et al. (2005). Measures of human population structure show heterogeneity among genomic regions. *Genome Research*, vol. 15, no. 11, pp. 1468-1476. doi:<http://dx.doi.org/10.1101/gr.4398405>

ZHAO, F. et al. (2015) Detection of selection signatures in dairy and beef cattle using high-density genomic information. *Genetic Selection Evolution*, vol. 47, pp. 49. doi:<http://dx.doi.org/10.1186/s12711-015-0127-3>