

Evaluation of Turkish Maize Landraces Through Observing Their Yield and Agro-Morphological Traits for Genetic Improvement of New Maize Cultivars

Ferhat Kizilgeci, Mehmet Yildirim, Onder Albayrak, Behiye Tuba Bicer, Akbar Hossain, Ayman EL Sabagh, Cuma Akinci

References

- AHMADI, V. et al. (2014) Correlation and path coefficient analyses of forage yield in corn hybrids as second crop. *Int. J. Biosci.*, vol. 4, no. 4, pp. 170-175. doi: <https://doi.org/10.12692/ijb/4.4.170-175>
- AKHTER, M.M. et al. (2016) Chlorophyll meter – a decision-making tool for nitrogen application in wheat under light soils. In *Int. J. Plant Prod.*, vol. 10, no. 3, pp. 289-302.
- AVLOV, J. et al. (2012) Relationship between grain yield, yield components and morphological traits in maize (*Zea mays* L.). In *Proceedings. 47th Croatian and 7th International Symposium on Agriculture*. Opatija.
- AZAR, C. et al. (1997). Maize landraces of the St. Lawrence-Great Lakes region of North America. *Euphytica*, vol. 98, no. 3, pp. 141-148.
- BOĆANSKI, J. et al. (2009) Genetic and phenotypic relationship between grain yield and components of grain yield of maize (*Zea mays* L.). *Genetika*, vol. 41, no. 2, pp. 145-154.
- BOSHEV, D. et al. (2014). Evaluation of maize hybrids for grain yield stability under rainfed and irrigated conditions using GGE biplot analysis. *Bulgarian J. Agric. Sci.*, vol. 20, no. 6, pp. 1320-1325.
- BRUSH, S. (1995) In situ conservation of landraces in centers of crop diversity. *Crop Sci.*, vol. 35, pp. 346–354.
- CARVALHO, I.R. et al. (2017) Components of variance and inter-relation of important traits for maize (*Zea mays*) breeding. *Aust. J. Crop Sci.*, vol. 11, no. 8, pp. 982-988. doi: <https://doi.org/10.21475/ajcs.17.11.08.pne474>
- De GALARRETA, J.R. and ALVAREZ, A. (2001) Morphological classification of maize landraces from northern Spain. *Genet. Resour. Crop Evol.*, vol. 48, no. 4, pp. 391-400.
- DEWEY, D.R. and LU, K.H. (1959) A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.*, vol. 51, pp. 515-518.
- ENCYCLOPAEDIA BRITANNICA. (2016) Zea. Website Name: *Encyclopædia Britannica*. <https://www.britannica.com/plant/Zea> (Accessed on May 05, 2018).
- ERDEN, I. (1991) *A research on determination of yield and yield characteristics of some hybrid and composite corn varieties in advanced generations (F1 and F2) in Samsun ecological conditions*: Master thesis. University of Ondokuz Mayıs.
- FERDOUSH, A. et al. (2017) Variability and traits association in maize (*Zea mays* L.) for yield and yield associated characters. In *J. Bangladesh Agric. Univ.*, vol. 15, no. 2, pp. 193-198.
- FETAHU S., et al. (2015) Genetic variability for yield and yield components among maize landraces. *ICAFE, Korçë Albania 25.09. 2015*, pp. 108-114.
- FRANKEL, O.H. et al. (1995) *The conservation of plant biodiversity*. Cambridge: Cambridge University Press.
- GOKMEN, S. (1995). Research on Yield and Yield Components of Hybrid and Composite Dent Corn Varieties in F1 and F2 Generations. *Turkish J. Agric. For.*, vol. 21, no. 3, pp. 267-272.
- GOKMEN, S. et al. (2001) Response of popcorn (*Zea mays* everta) to nitrogen rates and plant densities. *Turkish J. Agric. For.*, vol. 25 no. 1, pp. 15-23.

- GOODMAN, M.M. and PATERNIANI, E. (1969) The races of maize: III. Choices of appropriate characters for racial classification. *Econom. Bot.*, vol. 23 no. 3, pp. 265-273.
- GOZUBENLI, H. ET AL. (2001) THE EFFECT OF DIFFERENT NITROGEN DOSES ON GRAIN YIELD AND YIELD-RELATED CHARACTERS of some maize genotypes grown as second-crop. *J. Agric. Fac. CU.*, vol. 16, pp. 39-48.
- GOZUBENLI, H. et al. (2003) Effect of hybrid and plant density on grain yield and yield components of maize (*Zea mays*). In *Ind. J. Agron.*, vol., 48, pp. 203-205.
- HOQUE, M. M. et al. (2008) Genetic divergence in maize (*Zea mays* L.). *Bangladesh J. Agril. Res.*, vol., 9, pp.145–148
- IDIKUT, L. and KARA, S.N. (2011) The effects of previous plants and nitrogen rates on second crop corn. *Turkish J. Field Crops*, vol. 16, no. 2, pp. 239-244.
- KADIR, M.M. (2010) *Development of quality protein maize hybrids and their adaption in Bangladesh*: Ph.D. thesis. Mymensingh: Bangladesh Agricultural University, Department of Genetics and Plant Breeding.
- KHODARAHMPOUR, Z. (2012) Morphological Classification of Maize (*Zea mays* L.) Genotypes in Heat Stress Condition. *J. Agric. Sci.*, vol. 4, no. 5, pp. 43–76.
- KONUSKAN, O. (2000) *Effects of plant density on yield and yield-related characters of some maize hybrids grown in Hatay conditions as second crop*: M.Sc. Thesis. Thika: Mount Kenya University, Science Inst. MKU.
- KUMAR, A. et al. (2015) Diversity among maize landraces in North West Himalayan region of India assessed by agro-morphological and quality traits. *Ind. J. Genet. Plant Breed.*, vol. 75, no. 2, pp. 188-195.
- LANA, M.A. et al. (2017) Yield stability and lower susceptibility to abiotic stresses of improved open-pollinated and hybrid maize cultivars. *Agron. Sustain. Dev.*, vol. 37, pp. 30. doi: <https://doi.org/10.1007/s13593-017-0447-5>
- MENDES-MOREIRA, P. et al. (2015) Genetic Architecture of Ear Fasciation in Maize (*Zea mays*) under QTL Scrutiny. *PLoS ONE* vol. 10, no. 4: e0124543. doi: <https://doi.org/10.1371/journal.pone.0124543>
- MUCHIE, A. and FENTIE, D. (2016) Performance Evaluation of Maize Hybrids (*Zea Mays* L.) in Bahir Dar Zuria District, North Western Ethiopia, Department of natural sciences, Addis Zemen Preparatory school, Addis Zemen Ethiopia. *Intl. Res. J. Agric. Soil Sci.*, vol. 3, pp. 37–43.
- NZUVE, F. et al. (2013). Analysis of genotype x environment interaction for grain yield in maize hybrids. *J. Agric. Sci.*, vol. 5, no. 11, pp. 75-85. doi: <https://doi.org/10.5539/jas.v5n11p75>
- ONER, F. and GULUMSER, A. (2014) Determination of Some Agronomical Characteristics of Local Flint Corn (*Zea mays* L. *indurata*) Genotypes in The Black Sea Region of Turkey. *Türk Tarım ve Doğa Bilimleri*, vol. 7, no. 7, pp. 1800-1804.
- ONER, F. (2015) Determination of Chemical Quality Parameters with Yield and Yield components of Maize (*Zea mays* L.) Hybrids According to Various FAO Maturity Groups. In *J. Tekirdag Agric. Fac.*, vol. 12, no. 1, pp. 1-7
- PALUMBO, F. et al. (2017) Venetian Local Corn (*Zea mays* L.) Germplasm: Disclosing the Genetic Anatomy of Old Landraces Suited for Typical Cornmeal Mush Production. *Diversity*, vol. 9 no. 3, pp. 32.
- PAVAN, R. et al. (2011) Research Note Correlation and path coefficient analysis of grain yield and yield contributing traits in single cross hybrids of maize (*Zea mays* L.). *Electronic J. Plant Breed.*, vol. 2, no. 2, pp. 253-257.
- PEIFFER, J.A. et al. (2014) The genetic architecture of maize height. In *Genetics*, vol. 196, no. 4, pp. 1337-1356.
- RATHER, A.G. et al. (2003) Genetic variation in maize (*Zea mays* L.) population in high altitude temperate conditions in Kashmir. In *Indian J. Agril. Sci.*, vol. 79, no. 3, pp. 179–180.
- SANTOS, O.S.D. et al. (1993) Comparison of F1 and F2 Generations of Commercial Hybrids Maize. *Pcog. Agropec. Gros, Brasilia*, vol. 28, no. 1, pp. 75-79,

- SAS INSTITUTE INC. (1989). About: JMP 10 and Excel. Available at https://www.jmp.com/en_us/about.html (Accessed on 05 May 2018)
- SHANBAO, Q. et al. (2009) Effective improvement of genetic variation in maize lines derived from R08xDonor backcrosses by SSRs. *Biotech.*, vol. 8, pp.358–364.
- SHARIFI, R. S. et al. (2009) Effect of population density on yield and yield attributes of maize hybrids. *Res. J. Biol. Sci.*, vol. 4, no. 4, pp. 375-379.
- SHAW, R. H. (1988) Climate requirement. *corn and corn impr.*, pp. 609-638.
- TURGUT, I., (2000) Effects of Plant Populations and Nitrogen Doses on Fresh Ear Yield and Yield Components of Sweet Corn (*Zea mays saccharata* Sturt.) Grown Under Bursa Conditions. *Turk. J. Agric. For.*, vol. 24, pp. 341–347.
- TURGUT, I. et al. (2005) Alternate row spacing and plant density effects on forage and dry matter yield of maize hybrids (*Zea mays* L.). *J. Agron. Crop Sci.*, vol. 91, pp.146-151.
- TURKISH STATISTICAL INSTITUTE. (2017) *Turkish Statistical Institute: Statistics*. [Online]. Available from <http://www.tuik.gov.tr> [Accessed 2017-12-15]
- TUTEN C. and Demir I. (1984) Research on Yield and Yield Components in Advanced Generation of Hybrid and Composite Maize Varieties. *J. Agric. Fac. Ege Univ.*, pp.179-190.
- ULYSSES, J.G. (1963) Races of maize in Venezuela (Vol. 1136). Bogota: National Academy of Sciences.
- WASALA, S.K. et al. (2013). Analysis of yield performance and genotype x environment effects on selected maize (*Zea mays*) landrace accessions of India. *Ind. J. Agric. Sci.*, vol. 83, no.3, pp. 287-293.
- YILMAZ, S. et al. (2007) Genotype and Plant Density Effects on Corn (*Zea mays* L.) Forage Yield. *Asian J. Plant Sci.*, vol. 6, pp. 538-541. doi: <https://doi.org/10.3923/ajps.2007.538.541>
- ZHUKOVSKY, P.M. (1951). Ecological types and economic importance of Anatolian wheat (Translators: Kipçak, C., Nouruzhan, H. and Turkistanli, S.). pp. 158-214. *Turkish Sugar Beet Plants Publications: Agricultural Structure of Turkey* (in Turkish).
- ZSUZSANNA, Z. et al. (2002). Inheritance of plant and ear height in maize (*Zea mays* L.). *Acta Agraria Debreceniensis*, vol., 8, pp. 34-38.