

Comparative sensitivity of maize weevil to essential oil of *Hoslundia opposita* Vahl leaves subjected to different drying regimes

Samuel Adelani Babarinde, Oladele Abiodun Olaniran, Lamidi Ajao Usman, Ebenezer Oluwaseun Esan, Adeola Afolabi, Olawumi Sanmori, Joshua Damilare Lomowu

References

- AGAH, M. and NAJAFIAN, S. (2012) Essential oil content and composition of *Lippa citriodora* as affected by drying method before flowering stages. In *European Journal of Experimental Biology*, vol. 2, no 5, pp 1771-1777.
- ARABHOSSEINI, A. et al. (2006) Loss of essential oil of tarragon (*Artemisia dracunculus* L.) due to drying. In *Journal of the Science of Food and Agriculture*, vol. 86, pp 2543-2550. doi:<http://dx.doi.org/10.1002/jsfa.2641>
- BABARINDE G. O., BABARINDE S. A. and OGUNSOLA S. O. (2013a) Effect of maize weevil (*Sitophilus zeamais* Motschulsky 1855) infestation on the quality of three commercial pastas. In *Food Science and Quality Management*, vol. 21, pp 1-11.
- BABARINDE S. A. et al. (2013b) Effect of the prevalent insect species of yam chips on consumers' acceptability of yam paste. *Agriculturae Tropica Subtropica*, vol. 46, no. 3, pp 97-101. doi: <https://dx.doi.org/10.2478/ats-2013-0017>
- BABARINDE S. A. et al. (2008a) Potential of *Tithonia diversifolia* with Pirimiphos-methyl in control of *Sitophilus zeamais* (Coleoptera: Curculionidae). In *African Journal of Plant Science and Biotechnology*, vol. 2 no. 2, pp 77–80.
- BABARINDE S. A. et al. (2015) First report of toxicity of *Xylopiia parviflora* (A. Rich.) Benth (Annonaceae) root bark's essential oil against cowpea seed bruchid, *Callosobruchus maculatus* Fabricius (Coleoptera: Chrysomelidae: Bruchinae). In *Natural Product Research*, vol. 29 no. 4, pp 349-352. doi:<http://dx.doi.org/10.1080/14786419.2014.940943>
- BABARINDE S. A. et al. (2017a) Chemical composition of the essential oil of Nigeria grown *Hoslundia opposita* VAHL (Lamiaceae) dried leaves and its bioactivity against cowpea Seed Bruchid. In *Chemistry and Biodiversity*, vol.14, (In Press). doi:<https://dx.doi.org/10.1002/cbdv.201600418>
- BABARINDE S. A. et al. (2017b) Insectifugal and insecticidal potentials of two tropical botanical essential oils against cowpea seed bruchid. In *Environmental Science and Pollution Research* (In Press). doi:<https://dx.doi.org/10.1007/s11356-017-9589-x>
- BABARINDE S. A., SOSINA A. and OYEYIOLA E. I. (2008b) Susceptibility of selected crop in storage to *Sitophilus zeamais* Motschulsky in southwestern Nigeria. In *Journal of Plant Protection Research*, vol. 48, no. 4, pp 541-550.
- BABARINDE S. A. et al. (2014) Toxicity and repellence of *Hoslundia opposita* Vahl (Lamiaceae) leaves' essential oil against rust-red flour beetle, *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). In *Natural Product Research*, vol. 28, no. 6, pp 361-371. doi:<http://dx.doi.org/10.1080/14786419.2013.866115>
- BOUGHERRA H. H. et al. (2015) *Pistacia lentiscus* essential oil has repellent effect against three major insect pests of pasta. In *Industrial Crops and Products*, vol. 63, pp 249-255. doi:<http://dx.doi.org/10.1016/j.indcrop.2014.09.048>
- CONTI B. et al. (2010) Repellence of essential oils from tropical and Mediterranean Lamiaceae against *Sitophilus zeamais*. In *Bulletin of Insectology*, vol 63, no. 2, pp 197-202.
- HANAA, A. R. M. et al. (2012) Lemongrass (*Cymbopogon citratus*) essential oil as affected by drying methods. In *Annals of Agricultural Sciences*, vol. 57 no. 2, pp 113-116. doi: <http://dx.doi.org/10.1016/j.aogas.2012.08.004>

- HERRERA J. M. et al. (2014) Fumigant toxicity from five essential oils rich in ketones against *Sitophilus zeamais* (Motschulsky). In *Agriscientia*, vol. 31, no. 1, pp 35-41.
- KHALID K. A., HU W. and CAI W. (2008) The effects of harvesting and different drying methods on the essential oil composition of Lemon Balm (*Melissa officinalis* L.). In *Journal of Essential Oil Bearing Plants*, vol. 11, no. 4, pp 342-349. doi:<http://dx.doi.org/10.1080/0972060X.2008.10643639>
- KHANGHOLI S. P. and REZAEINODEHI A. (2008) Effect of drying temperature on essential oil content and composition of Sweet Wonnwood (*Artemisia annua*) growing wild in Iran. In *Pakistan Journal of Biological Sciences*, vol. 11, no. 6, pp 934-937.
- Lira C. S. et al. (2015) Evaluation of the toxicity of essential oil from *Alpinia purpurata* inflorescences to *Sitophilus zeamais* (maize weevil). In *Crop Protection*, vol. 71, pp 95-100. Doi: 10.1016/j.cropro.2015.02.004
- MAHESWARAN R. and IGNACIMUTHU S. (2013) Bioefficacy of essential oil from *Polygonum hydropiper* L. against mosquitoes, *Anopheles stephensi* and *Culex quinquefasciatus*. In *Ecotoxicology and Environmental Safety*, vol. 97 no. 1, pp 26-31. doi:<https://dx.doi.org/10.1016/j.ecoenv.2013.06.028>
- MOHARRAMIPOUR S. and NEGAHBAN M. (2014) Plant essential oils and pest management. In: SAHAYARAJ, K. (ed): *Basic and applied aspects of biopesticides*. New Delhi: Springer, India. doi:<http://dx.doi.org/10.1007/978-81-322-1877-7-7>
- OMIDBAIGI R., SEFIDKON F. and KAZEMI F. (2004) Influence of drying methods on the essential oil content and composition of Roman chamomile. In *Flavour and Fragrance Journal*, vol. 19, no. 3, pp 196-198. <http://dx.doi.org/10.1002/ffj.1340>
- PIMIENTA-RAMÍREZ L. et al. (2016) Chemical composition and evaluation of the essential oil from *Eupatorium glabratum* as biopesticide against *Sitophilus zeamais* and several stored maize fungi. In *Journal of Essential Oil Research*, vol. 28, no. 2, pp 113-120. doi:<http://dx.doi.org/10.1080/10412905.2015.1093969>
- RAHIMMALEK M. and GOLI S. A. H. (2013) Evaluation of six drying treatments with respect to essential oil yield, composition and color characteristics of *Thymus daenensis* subsp. *daenensis*. Celak leaves. In *Industrial Crops and Products*, vol. 42, pp 613-619. doi:<http://dx.doi.org/10.1016/j.indcrop.2012.06.012>
- SANTOS A. C. V. et al. (2015) Use of plant oils from the southwestern Amazon for the control of maize weevil. In *Journal of Stored Product Research*, vol. 63, pp 67-70. doi:<http://dx.doi.org/10.1016/j.jspr.2015.07.002>
- SELLAMI I. H. et al. (2011). Qualitative and quantitative changes in the essential oil of *Laurus nobilis* L. leaves as affected by different drying methods. In *Food Chemistry*, vol. 126, no. 2, pp 691-697. doi:<http://dx.doi.org/10.1016/j.foodchem.2010.11.022>
- SHAHHOSEINI R. et al. (2013) The effect of different drying methods on the content and chemical composition of essential oil of Lemon verbena (*Lippia citriodora*). In *Journal of Essential Oil Bearing Plants*, vol. 16, No. 4, pp 474–81. doi:<http://dx.doi.org/10.1080/0972060X.2013.813270>
- SPSS. (2006) *Statistical Package for Social Sciences. Version 15.0 for Windows*.
- TELES S. et al. (2013). Effect of geographical origin on the essential oil content and composition of fresh and dried *Mentha villosa* Hudson leaves. In *Industrial Crops and Products*, vol. 46, no. 1; pp 1-7. doi:<http://dx.doi.org/10.1016/j.indcrop.2012.12.009>
- USMAN L. A. et al. (2016) Phytochemical profile and insecticidal activity of essential oil from fresh and dried leaves of the Nigerian grown *Citrus meyeri*. In *Journal of Turkish Chemical Society (Section A)*, vol. 3, no. 2, pp 207-218. doi:<http://dx.doi.org/10.18596/jotcsa.09331>