

References

- Adeyemi, N.O. et al. (2020). Effect of commercial arbuscular mycorrhizal fungi inoculant on growth and yield of soybean under controlled and natural field conditions, *Journal of Plant Nutrition*, 43:4, 487-499, DOI: 10.1080/01904167.2019.1685101
- Adeyemi, N.O. et al. (2019). Identification and relative abundance of native arbuscular mycorrhizal fungi associated with oil-seed crops and maize (*Zea mays* L.) in derived savannah of Nigeria. In *Acta fytotechn zootech*, 22, 2019(3): 84–89
- Adeyemi, N., Sakariyawo, O. and Atayese, M., (2017). Yield and yield attributes responses of soybean (*Glycine max* L. Merrill) to elevated CO₂ and arbuscular mycorrhizal fungi inoculation in the humid transitory rainforest. *Notulae Scientia Biologicae*, 9(2):233-241. DOI: 10.15835/nsb9210002.
- Ainsworth, E. A., and Rogers, A. (2007). The response of photosynthesis and stomatal conductance to rising [CO₂]: Mechanisms and environmental interactions. *Plant, Cell and Environment*, 30, 258–270. doi:10.1111/j.1365- 3040.2007.01641.x
- Alberton, O., Kuyper, T.W., and Gorissen, A., (2005). Taking mycocentrism seriously: mycorrhizal fungal and plant responses to elevated CO₂. *NewPhytol.*167, 859–868
- Bardgett, R.D., Freeman, C., and Ostle, N.J., (2008). Microbial contributions to climate change through carbon cycle feedbacks. *ISME J.* 2, 805-814.
- Bécard, G. et al. (1992). Extensive in vitro hyphal growth of vesicular-arbuscular mycorrhizal fungi in presence of CO₂ and flavenols. *Applied Environmental Microbiology* 58, 821–825.
- Bhattacharyya, P., et al. (2016). Elucidation of rice rhizosphere metagenome in relation to methane and nitrogen metabolism under elevated carbon dioxide and temperature using whole genome metagenomic approach. *Sci. Total Environ* 542, 886–898.
- Brundrett M.C. (2009). Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil* 320: 37–77.
- Butterly C.R. et al. (2016). Long-term effects of elevated CO₂ on carbon and nitrogen functional capacity of microbial communities in three contrasting soils. *Soil Biology and Biochemistry* 97, 157-167
- Cairney, J.W.G. (2012). Extramatrical mycelia of ectomycorrhizal fungi as moderators of carbon dynamics in forest soil. *Soil Biology and Biochemistry* 95, 198-208.
- Cannell, M.G.R. and Thornley, H.M. (1998). N-poor ecosystems may respond more to elevated [CO₂] than N-rich ones in the long term. A model analysis of grassland. *Glob. Change Biol.* 4, 431–442.
- Carrillo, Y. et al. (2014). Plant rhizosphere influence on microbial C metabolism: the role of elevated CO₂, N availability and root stoichiometry. *Biogeochemistry* 117, 229-240.

- Cheng, L. et al. (2011). Soil microbial responses to elevated CO₂ and O₃ in a nitrogen-aggrading agroecosystem. *Plos One* 6, e21377.
- Compant, S. et al. (2010). Climate change effects on beneficial plant–microorganism interactions. *Microbiology Ecology* 73, 197–214.
- Cotton, T.E. et al. (2015). Fungi in the future: interannual variation and effects of atmospheric change on arbuscular mycorrhizal fungal communities. *New Phytol.* 205 (4), 1598–1607.
- Denef, K. et al. (2007). Community shifts and carbon translocation within metabolically-active rhizosphere microorganisms in grasslands under elevated CO₂. *Biogeosciences* 4, 769-779
- Drigo, B. et al. (2013). Impacts of 3 years of elevated atmospheric CO₂ on rhizosphere carbon flow and microbial community dynamics. *Glob. Change Biol.* 19, 621-636.
- Drissner, D. et al. (2007). Nine years of enriched CO₂ changes the function and structural diversity of soil microorganisms in a Grassland. *Eur. J. Soil Sci.* 58, 260-269.
- Fang, H.J. et al. (2015). Elevated atmospheric carbon dioxide concentration stimulates soil microbial activity and impacts water-extractable organic carbon in an agricultural soil. *Biogeochemistry* 122, 253-267.
- Gavito M.E. et al. (2000) Atmospheric CO₂ and mycorrhiza effects on biomass allocation and nutrient uptake of nodulated pea (*Pisum sativum* L.) plants. *J Exp Bot*; 51:1931–8.
- Ghannoum, O. et al., (2010). Exposure to preindustrial, current and future atmospheric CO₂ and temperature differentially affects growth and photosynthesis in Eucalyptus. *Global Change Biology* 16, 303-319.
- Giovanetti, M. and Mosse, B. (1980). An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phytologist* 84, 489–500.
- Goicoechea, N. et al. (2014) Increased photosynthetic acclimation in alfalfa associated with arbuscular mycorrhizal fungi (AMF) and cultivated in greenhouse under elevated CO₂ *Journal of Plant Physiology* 171: 1774–1781
- Haugwitz, M.S. et al. (2014). Soil microorganisms respond to five years of climate change manipulations and elevated atmospheric CO₂ in a temperate heath ecosystem. *Plant Soil* 374, 211-222.
- Huang, X. et al. (2014). Changes of soil microbial biomass carbon and community composition through mixing nitrogen–fixing species with *Eucalyptus urophylla* in subtropical China. *Soil Biol. Biochem.* 73, 42–48.
- Ineichen, K. Wiemken, V. and Wiemken, A. (1995). Shoots, roots and ectomycorrhiza formation of pine seedlings at elevated atmospheric carbon dioxide. *Plant Cell Environ.* 18, 703–707.
- IPCC (2013). “Summary for Policymakers,” in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*

Intergovernmental Panel on Climate Change, (eds) T.F.Stocker, D .Qin,G.K. Plattner, M. Tignor, S.K. Allen ,J. Boschung.(NewYork, NY: Cambridge UniversityPress).

Jin, J. et al. (2013). Elevated CO₂ temporally enhances phosphorus immobilization in the rhizosphere of wheat and chickpea. *Plant Soil* 368, 315-328

Johnson, N.C. et al. (2013). Predicting community and ecosystem outcomes of mycorrhizal responses to global change. *Ecology Letters* 16, 140-153.

Johnson, N.C. et al. (2005). Species of plants and associated arbuscular mycorrhizal fungi mediate mycorrhizal responses to CO₂ enrichment. *Global Change Biology* 11, 1156-1166.

Johnson, N.C. and Gehring, C.A. (2007). Mycorrhizas: symbiotic mediators of rhizosphere and ecosystem processes. In: Cardon, Z.G., Whitbeck, J.L. (Eds.), *The Rhizosphere: An Ecological Perspective*. Elsevier Academic Press, London, UK, pp. 31-56

Kabir Z. et al. (1997). Seasonal changes of arbuscular mycorrhizal fungi as affected by tillage practices and fertilization: hyphal density and mycorrhizal root colonization. *Plant Soil*. 192(2):285–293. doi:10.1023/A:1004205828485.

Kumar, A. et al. (2019). Effects of water deficit stress on agronomic and physiological responses of rice and greenhouse gas emission from rice soil under elevated atmospheric CO₂. *Sci. Total Environ.* 650, 2032–2050

Kuzyakov, Y. et al. (2018) Review and synthesis of the effects of elevated atmospheric CO₂ on soil processes: No changes in pools, but increased fluxes and accelerated cycles, *Soil Biology and Biochemistry*, doi: 10.1016/j.soilbio.2018.10.005

Liu, S., et al. (2018). Climatic role of terrestrial ecosystem under elevated CO₂: a bottom-up greenhouse gases budget. *Ecology Letters* 21, 1108-1118.

Matamala, R. and Drake, B.G. (1999). The influence of atmospheric CO₂ enrichment on plant–soil nitrogen interactions in a wetland plant community on the Chesapeake Bay. *Plant Soil* 210, 93–101

Mccarthy, H. R. et al. (2010). Re-assessment of plant carbon dynamics at the Duke free-air CO₂ enrichment site: Interactions of atmospheric CO₂ with nitrogen and water availability over stand development. *New Phytologist* 185 (2):514–28. doi: 10.1111/j.1469-8137.2009.03078.x.

Moran, K.K. and Jastrow, J.D. (2010). Elevated carbon dioxide does not offset loss of soil carbon from a corn–soybean agroecosystem. *Environmental Pollution* 158, 1088–1094.

Nie, M. et al. (2013). Positive climate feedbacks of soil microbial communities in a semi-arid grassland. *Ecol. Lett.* 16, 234-241

Oliveira V.F. et al (2010). Elevated CO₂ atmosphere promotes plants growth and inulin production in the cerrado species *Vernonia herbacea*. *Funct Plant Biol*; 37:223–31

- Panneerselvam P. et al. (2019) Influence of elevated CO₂ on arbuscular mycorrhizal fungal community elucidated using Illumina MiSeq platform in sub-humid tropical paddy soil. *Applied Soil Ecology* 145; 103344, doi.org/10.1016/j.apsoil.2019.08.006
- Pendall, E. et al. (2013). Warming reduces carbon losses from grassland exposed to elevated atmospheric carbon dioxide. *PLoS One* 8, e71921.
- Phillips, J.M. and Hayman D.S. (1970). Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. British Mycology Society* 55, 150–160.
- Procter, A.C. et al. (2014). Fungal community responses to past and future atmospheric CO₂ differ by soil type. *Appl. Environ. Microbiol.* 80, 7364-7377.
- Reinsch, S. et al. (2013). Impact of future climatic conditions on the potential for soil organic matter priming. *Soil Biol. Biochem.* 65, 133-140.
- Rillig, M.C. and Allen, M.F. (1999). What is the role of arbuscular mycorrhizal fungi in plant-to-ecosystem responses to elevated atmospheric CO₂? *Mycorrhiza* 9,1–8.
- Rogers A, Y. et al, (2006). Increased C availability at elevated carbon dioxide concentration improves N assimilation in a legume. *Plant, Cell and Environment*: 29, 1651-1658.
- Saitoh Y, et al. (2004). Yeast generated CO₂ as a convenient source of CO₂ for adult mosquito sampling. *Journal of American Mosquitoes Control Association* 20:261–264
- Sakariyawo O.S. et al. (2016) Growth, assimilate partitioning and grain yield response of soybean (*Glycine max* L. Merrill) varieties to carbon dioxide enrichment and arbuscular mycorrhizal fungi in the humid rainforest. *Agro-science*: 15, 29-40
- Schortemeyer, M. et al. (2000). Effects of elevated atmospheric CO₂ concentration in C and N pools and rhizosphere processes in a Florida scrub oak community. *Glob. Change Biol.* 6, 383–391.
- Sillen, W.M.A. and Dieleman W.I.J. (2012). Effects of elevated CO₂ and N fertilization on plant and soil carbon pools of managed grasslands: a meta-analysis. *Biogeosciences* 9, 2247-2258.
- Singh, S.B., et al. (2017). Impact of secondary forest fallow period on soil microbial biomass carbon and enzyme activity dynamics under shifting cultivation in North Eastern Hill region, India. *Catena* 156, 10–17.
- Singh, L.P. Gill, S.S. and Tuteja, N. (2011). Unravelling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Signal. Behav.* 6 (2), 175–191.
- Smith, S.E. and Read, D.J., (2008). *Mycorrhizal Symbiosis*, third ed. Academic Press (Elsevier Ltd.), London, UK, pp. 316e319.
- Vance E.D. (1987): An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.*, 19, 703–707. doi:10.1016/0038-0717(87) 90052-6

Weigel, H.J. et al. (2005). Carbon turnover in a crop rotation under free air CO₂ enrichment (FACE). *Pedosphere* 15, 728-738.

Wipf, D. et al. (2019). Trading on the arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks. *New Phytol.* 223 (3), 1127–1142.

Wu, D.X. et al. (2004). Effects of elevated CO₂ concentration on growth, water use, yield and grain quality of wheat under two soil water levels. *Agriculture, Ecosystems & Environment* 104, 493–507.

Yang, L. et al. (2006). Seasonal changes in the effects of free-air CO₂ enrichment (FACE) on dry matter production and distribution of rice (*Oryza sativa* L.). *Field Crops Research*, 98, 12–19. doi:10.1016/j.fcr.2005.11.003

Ziska, L. H. (2000). The impact of elevated CO₂ on yield loss from a C₃ and C₄ weed in field-grown soybean. *Global Change Biology* 6:899–905.