

The effect of natural feed additive on productive performance of broiler chickens

Cyril Hrnčár, Martin Fik, Emília Hanusová, Anton Hanus, Henrieta Arpášová, Marie Hamadová, Tomáš Kanka, Jozef Bujko

References

- Abou-Elkhair, R. et al. (2014). Effects of black pepper (*Piper nigrum*), turmeric powder (*Curcuma longa*) and coriander seeds (*Coriandrum sativum*) and their combinations as feed additives on growth performance, carcass traits, some blood parameters and humoral immune response of broiler chickens. *Asian-Australasian Journal of Animal Sciences*, 27, 847–854. [doi: 10.5713/ajas.2013.13644](https://doi.org/10.5713/ajas.2013.13644)
- Akbarian, A. et al. (2012). Influence of turmeric rhizome and black pepper on blood constituents and performance of broiler chickens. *African Journal of Biotechnology*, 11(34), 8606–8611. [doi: 10.5897/AJB11.3318](https://doi.org/10.5897/AJB11.3318)
- Al-Harhi, M.A. (2006). Impact of supplemental feed enzymes, condiments mixture or their combination on broiler performance, nutrients digestibility and plasma constituents. *International Journal of Poultry Science*, 5(8), 764–771. [doi: 10.3923/ijps.2006.764.771](https://doi.org/10.3923/ijps.2006.764.771)
- Ali, A.H.H. (2014). Productive performance and immune response of broiler chicks as affected by dietary marjoram leaves powder. *Egyptian Poultry Science Journal*, 34, 57–70. [doi: 10.21608/epsj.2014.5306](https://doi.org/10.21608/epsj.2014.5306)
- Al-Jaleel, R. & Abd A. (2012). Use of tumeric (*Curcuma longa*) on the performance and some physiological traits on the broiler diets. *The Iraqi Journal of Veterinary Medicine*, 36(1), 51–57.
- Al-Sultan, S.I. (2003). The effects of *Curcuma longa* (turmeric) on overall performance of broiler chickens. *International Journal of Poultry Science*, 2(5), 351–353.
- Attia, Y.A. et al. (2014). Productive performance, biochemical and hematological traits of broiler chickens supplemented with propolis, bee pollen, and mannan oligosaccharides continuously or intermittently. *Livestock Science*, 164, 87–95. <https://doi.org/10.1016/j.livsci.2014.03.005>
- Brugalli, I. (2003) Alimentacao alternativa: An utilizacao de fitoterapicos ou nutraceuticos como moduladores da imunidade e desempenho animal. *Anais do Simposio sobre Manejo e Nutricao de Aves e Suínos*, Campinas, Sao Paulo. Brasil. Campinas: CBNA, 167–182.
- Capcarová, M. et al. (2010). Effect of *Lactobacillus fermentum* and *Enterococcus faecium* strains on internal milieu, antioxidant status and body weight of broiler chickens. *Journal of Animal Physiology and Animal Nutrition*, 94(5), 215–224. [doi: 10.1111/j.1439-0396.2010.01010.x](https://doi.org/10.1111/j.1439-0396.2010.01010.x)
- Capcarová, M. et al. (2014). Comparative analysis of acetic and citric acid on internal milieu of broiler chickens. *Potravinarstvo Slovak Journal of Food Sciences*, 8(1), 190–195. [doi: 10.5219/379](https://doi.org/10.5219/379)
- Daneshyar, A. et al. (2012). The effects of turmeric supplementation on antioxidant status, blood gas indices and mortality in broiler chickens with T(3)-induced ascites. *British Poultry Science*, 53(3), 379–385. [doi: 10.1080/00071668.2012.702340](https://doi.org/10.1080/00071668.2012.702340)
- Daneshmand, A. et al. (2015). Evaluating complementary effects of ethanol extract of propolis with the probiotic on growth performance, immune response and serum metabolites in male broiler chickens. *Livestock Science*, 178, 195–201. <https://doi.org/10.1016/j.livsci.2015.04.012>
- Doley, S. et al. (2009). Effect of supplementation of ginger, garlic and turmeric in broiler chicken. *Indian Veterinary Journal*, 86(6), 644–645.
- Duncan, D. B. (1955). Multiple ranges and multiple F-test. *Biometric*, 11, 10–42. [doi:10.2307/3001478](https://doi.org/10.2307/3001478)
- Durrani, F.R. et al. (2006). Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agriculture and Biological Sciences*, 1, 9–11.
- Emadi, M. & Kermanshahi, H. (2006). Effect of turmeric rhizome powder on performance and carcass characteristics of broiler chickens. *International Journal of Poultry Science*, 5(11), 1069–1072. [doi: 10.3923/ijps.2006.1069.1072](https://doi.org/10.3923/ijps.2006.1069.1072)
- Fernandez, S. R. et al. (1994). Limiting order of amino acids in corn and soybean meal for growth of the chick. *Poultry Science*, 73, 1887–1896.
- Gheisari, A. et al. (2017). Effect of ethanolic extract of propolis as an alternative to antibiotics as a growth promoter on broiler performance, serum biochemistry, and immune responses. *Veterinary World*, 10(2), 249–254. [doi: 10.14202/vetworld.2017.249-254](https://doi.org/10.14202/vetworld.2017.249-254)
- Hashemi, S.R. and Davoodi, H. (2010). Phyto-genics as new class of feed additive in poultry industry. *Journal of Animal and Veterinary Advances*, 9, 2295–2304. [doi: 10.3923/javaa.2010.2295.2304](https://doi.org/10.3923/javaa.2010.2295.2304)

- Haščík, P. et al. (2019). The effect of bee products and probiotic on meat performance of broiler chickens. *Journal of Microbiology, Biotechnology and Food Sciences*, 9 (1), s. 88–92.
- Haščík, P. et al. (2020). The amino acid profile of broiler chicken meat after dietary administration of bee products and probiotics. *Biologia*, 75, 1–10.
- Hernández, F. et al. (2004). Influence of two plant extracts on broilers performance, digestibility, and digestive organ size. *Poultry Science*. 83, 169–174.
- Hrnčár, C. et al. (2015). Effect of L-carnitine supplementation on fattening and carcass parameters of broiler chickens. *Acta fytotechnica et zootecnica*, 18(1), 15–19. [doi: 10.15414/afz.2015.18.01.15–19](https://doi.org/10.15414/afz.2015.18.01.15-19)
- Hrnčár, C. et al. (2016) Effect of three-strains probiotic on productive performance and carcass characteristics of broiler chickens. *Scientific Papers: Animal Science and Biotechnologies*, 49(2), 149–154.
- Hrnčár, C. & Bujko, J. (2017). Effect of different levels of green tea (*Camellia sinensis*) on productive performance, carcass characteristics and organs of broiler chickens. *Potravinárstvo Slovak Journal of Food Sciences*, 11(1), 623–628. [doi: https://dx.doi.org/10.5219/809](https://dx.doi.org/10.5219/809)
- Hrnčár, C. et al. (2017). Effect of adding L-carnitine and probiotic on performance and carcass parameters of broiler chickens. *Scientific Papers: Animal Science and Biotechnologies*, 50(2), 86-93.7
- Hrnčár, C. et al. (2018). The effect of single and combined use of probiotic and humate on fattening performance, carcass characteristics and internal organs of broiler chickens. *Macedonian Journal of Animal Science*, 8(2), 81–87.
- Hui, L. et al. (2010). Chemical composition of lavender essential oil and its antioxidant activity and inhibition against rhinitisrelated bacteria. *African Journal of Microbiology Research*, 4 (4), 309–313.
- Hussein, S. N. (2013). Effect of turmeric (*Curcuma longa*) powder on growth performance, carcass traits, meat quality and serum biochemical parameters in broilers. *Journal of Advanced Biomedical & Pathobiology Research*, 3(2), 25–32.
- INTERMAG (2015). Available on <https://intermag.eu/livestock-production/product/musculaton>
- Jamroz, D. et al. (2003). The influence of phytogetic extracts on performance nutrient digestibility, carcass characteristics, and gut microbial status in broiler chickens. *Journal of Animal and Feed Sciences*, 12(3), 583–596. [doi: https://doi.org/10.22358/jafs/67752/2003](https://doi.org/10.22358/jafs/67752/2003)
- JASP 0.8.6 software (2018). Available on <https://jasp-stats.org/>
- Kalařová, A. et al. (2014). Metabolic effect of citric acid in broiler chickens. *Journal of Microbiology, Biotechnology and Food Sciences*, 3 (special issue), 110–112.
- Kopecký, J. et al. (2012). Effect of organic acids supplement on performance of broiler chickens. *Scientific Papers: Animal Science and Biotechnologies*, 2012, 45(1), 51–54.
- Král, M. et al. (2012). Application of probiotics in poultry production. *Scientific Papers. Animal Science and Biotechnologies*, 45, 55–57.
- Lambert, R.J.W. et al. (2001). A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology*, 91, 453-462. [doi: 10.1046/j.1365-2672.2001.01428.x](https://doi.org/10.1046/j.1365-2672.2001.01428.x).
- Mehala, C. & Moorthy, M. (2008). Effect of Aloe vera and *Curcuma longa* (turmeric) on carcass characteristics and biochemical parameters of broilers. *International Journal of Poultry Science*, 7, 857–861. [doi:10.3923/ijps.2008.857.861](https://doi.org/10.3923/ijps.2008.857.861)
- Murugesan, G.R. et al. (2015). Phytogetic feed additives as an alternative to antibiotic growth promoters in broiler chickens. *Frontiers of Veterinary Science*, 2(21). [doi: 10.3389/fvets.2015.00021](https://doi.org/10.3389/fvets.2015.00021)
- Nouzarian, R. et al. (2011). Effect of turmeric powder on performance, carcass traits, humoral immune responses, and serum metabolites in broiler chickens. *Journal of Animal and Feed Sciences*, 20, 389–400. [doi:10.22358/jafs/66194/2011](https://doi.org/10.22358/jafs/66194/2011)
- Ocak, N. et al. (2008). Performance of broilers fed diets supplemented with dry peppermint (*Mentha piperita* L.) or thyme (*Thymus vulgaris* L.) leaves as growth promoter source. *Czech Journal of Animal Science*, 53 (4), 169–175. [doi: 10.17221/373-CJAS](https://doi.org/10.17221/373-CJAS)
- Olukosi, O. A. & Dono, N. D. (2014). Modification of digesta pH and intestinal morphology with the use of benzoic acid or phytobiotics and the effects on broiler chicken growth performance and energy and nutrient utilization. *Journal of Animal Science*, 92, 3945–3953. [doi:10.2527/jas.2013-6368](https://doi.org/10.2527/jas.2013-6368)
- Ojano-Dirain, C. & Waldroup, P. (2002). Evaluation of lysine, methionine and threonine needs of broilers three to six week of age under moderate temperature stress. *International Journal of Poultry Science*, 1, 16–21. [doi:10.3923/ijps.2002.16.21](https://doi.org/10.3923/ijps.2002.16.21)

- Osman, M. et al. (2010). Productive, physiological, immunological and economical effects of supplementing natural feed additives to broiler diets. *Egyptian Poultry Science Journal*, 30(1), 25–53.
- Pistová, V. et al. (2016). The effect of the humic acid and garlic (*Allium sativum* L.) on performance parameters and carcass characteristic of broiler chicken. *Journal of Central European Agriculture*, 17(4), 1168–1178. [doi: 10.5513/JCEA01/17.4.1826](https://doi.org/10.5513/JCEA01/17.4.1826)
- Roofchae, A. et al. (2011). Effect of dietary oregano (*Origanum vulgare* L.) essential oil on growth performance, cecal microflora and serum antioxidant activity of broiler chickens. *African Journal of Biotechnology*, 10, 6177–6183. [DOI:10.4314/AJB.V10I32](https://doi.org/10.4314/AJB.V10I32)
- Song R.I. et al. (2017). Effects of dietary oregano powder supplementation on the growth performance, antioxidant status and meat quality of broiler chicks. *Italian Journal of Animal Science*, 16 (2), 246–252. <https://doi.org/10.1080/1828051X.2016.1274243>
- Sugiharto, S. (2016). Role of nutraceuticals in gut health and growth performance of poultry. *Journal of the Saudi Society of Agricultural Sciences*, 15(2), 99–111. [doi:10.1016/j.jssas.2014.06.001](https://doi.org/10.1016/j.jssas.2014.06.001)
- Suriya, R. et al. (2012). The effect of dietary inclusion of herbs as growth promoter in broiler chickens. *Journal of Animal and Veterinary Advances*, 11, 346–350. [doi:10.3923/javaa.2012.346.350](https://doi.org/10.3923/javaa.2012.346.350)
- Tesseraud, S. et al. M. (1996). Relative responses of protein turnover in three different skeletal muscles to dietary lysine deficiency in chicks. *British Poultry Science*, 37, 641–650. [doi:10.1080/00071669608417893](https://doi.org/10.1080/00071669608417893)
- Ultee, A. et al. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and Environmental Microbiology*, 68, 1561–1568. [doi: 10.1128/AEM.68.4.1561-1568.2002](https://doi.org/10.1128/AEM.68.4.1561-1568.2002)
- Wang, D. et al. (2015). Effects of dietary supplementation with turmeric rhizome extract on growth performance, carcass characteristics, antioxidant capability, and meat quality of Wenchang broiler chickens. *Italian Journal of Animal Science*, 14(3), 344–349. <https://doi.org/10.4081/ijas.2015.3870>
- Weis, J. et al. (2010) Effect of probiotic strain *Lactobacillus fermentum* CCM 7158 supplement on performance and carcass characteristics of broiler chickens. *Acta fytotechnica et zootechnica*, 13(4), 96–98.
- Zafarnejad, K. et al. (2017). Effect of bee glue on growth performance and immune response of broiler chickens. *Journal of Applied Animal Research*, 45(1), 280–284. <https://doi.org/10.1080/09712119.2016.1174130>