

## Candidate genes for congenital malformations in pigs

Zuzana Krupová, Emil Krupa, Eliška Žáková, Ludmila Zavadilová, Eva Kašná

### References

- Atkinson M. et al. (2017). Evaluation of the effect of umbilical hernias on play behaviors in growing pigs. *Canadian Veterinary Journal*, 58, 1065–1072.
- Bourneuf E. et al. (2018). New susceptibility loci for cutaneous melanoma risk and progression revealed using a porcine model. *Oncotarget*, 9(45), 27682–27697. <https://www.oncotarget.com/article/25455/text/>
- Brening B. et al. (2015). Porcine SOX9 gene expression is influenced by an 18bp indel in the 5'-untranslated region. *PLoS ONE*, 10(10), e0139583. <https://doi.org/10.1371/journal.pone.0139583>
- Cassini P. et al. (2005). Genetic analysis of anal atresia in pigs: evidence for segregation at two main loci. *Mammalian Genome*, 16(10), 164–170. <https://doi.org/10.1007/s00335-004-3024-6>
- Ding N.S. et al. (2009). A genome-wide scan reveals candidate susceptibility loci for pig hernias in an intercross between white Duroc and Erhualian. *Journal of Animal Science*, 87(8), 2469–2474. <https://doi.org/10.2527/jas.2008-1601>
- Du Z.Q. et al. (2009). Association and haplotype analyses of positional candidate genes in five genomic regions linked to scrotal hernia in commercial pig lines. *PLoS One*, 4(3), 4837. <https://doi.org/10.1371/journal.pone.0004837>
- Grindflek E. et al. (2018). Genome-wide association study reveals a QTL and strong candidate genes for umbilical hernia in pigs on SSC14. *Genomics*, 19, 412. <https://doi.org/10.1186/s12864-018-4812-9>
- Grindflek E. et al. (2006). Genome-wide linkage analysis of inguinal hernia in pigs using affected sib pairs. *BMC Genetics*, 7, 25. <https://doi.org/10.1186/1471-2156-7-25>
- Hao X. et al. (2017). Genome-wide association study identifies candidate genes for piglet splay leg syndrome in different populations. *BMC Genetics*, 18, 64. <https://doi.org/10.1186/s12863-017-0532-4>
- Hori T. et al. (2001). Mapping loci causing susceptibility to anal atresia in pigs, using a resource pedigree. *Journal of Pediatric Surgery*, 36, 1370–1374. <https://doi.org/10.1053/jpsu.2001.26373>
- Jin Q. et al. (2013). Molecular characterization and genome-wide mutations in porcine anal atresia candidate gene GLI2. *Mammalian Genome*, 24, 500–507. <https://link.springer.com/article/10.1007%2Fs00335-013-9485-8>
- Krupová Z. et al. (2017). New breeding objectives for the Czech pig population. *Indian Journal of Animal Sciences*, 87(6), 778–781.
- Liao X. et al. (2015). Susceptibility loci for umbilical hernia in swine detected by genome-wide association. *Russian Journal of Genetics*, 51(10), 1000–1006. <https://doi.org/10.1134/S1022795415100105>
- Lindholm-Perry A.K. et al. (2010). Genomic regions associated with kyphosis in swine. *BMC Genetics*, 11, 112. <https://doi.org/10.1186/1471-2156-11-112>
- Long Y. et al. (2016). A genome-wide association study of copy number variations with umbilical hernia in swine. *Animal Genetics*, 47, 298–305. <https://doi.org/10.1111/age.12402>
- Mahmud M.A. et al. (2015). Cryptorchidism in mammals - a review. *Global Journal of Animal Scientific Research*, 3(1), 128–135.
- Mattsson P. (2011). Prevalence of congenital defects in Swedish Hampshire, Landrace and Yorkshire pig breeds and opinions on their prevalence in Swedish commercial herds (MSc Thesis). Sweden, Uppsala: Swedish University of Agricultural Sciences, 35 p.
- Maak S. et al. (2009). Identification of candidate genes for congenital splay leg in piglets by alternative analysis of DNA microarray data. *International Journal of Biological Sciences*, 5(4), 331–337. <https://www.ijbs.com/v05p0331.htm>
- Misdorp W. (2003). Congenital and hereditary tumours in domestic animals. 2. Pigs. A review. *Veterinary Quarterly*, 25(1), 17–30. <https://doi.org/10.1080/01652176.2003.9695141>
- Moravčíková N. and Kasarda R. (2020). Use of High-density SNP analyses to develop a long-term strategy for conventional populations to prevent loss of diversity – review. *Acta fytotechnica et zootechnica*, 23, 236–240. <https://doi.org/10.15414/afz.2020.23.04.236-240>
- Morey-Matamalas A. et al. (2021). Neoplastic lesions in domestic pigs detected at slaughter: literature review and a 20-year review (1998–2018) of carcass inspection in Catalonia. *Porcine Health Management*, 7, 30. <https://doi.org/10.1186/s40813-021-00207-0>

- Nowacka-Woszuk J. (2021). The genetic background of hernia in pigs: A review. *Livestock Science*, 244, 104317. <https://doi.org/10.1016/j.livsci.2020.104317>
- Papatsiros V.G. (2012). The splay leg syndrome in piglets: a review. *American Journal of Animal and Veterinary Sciences*, 7(2), 80–83.
- Prunier A. et al. (2009). High physiological demands in intensively raised pigs: impact on health and welfare. *Animal*, 4(6), 886–898. <https://doi.org/10.1017/S175173111000008X>
- Qiushi J. et al. (2013). Molecular characterisation and genome-wide mutations in porcine anal atresia candidate gene GLI2. *Mammalian Genome*, 24, 500–507. <https://doi.org/10.1007/s00335-013-9485-8>
- Rohrer G.A. et al. (2015). A study of vertebra number in pigs confirms the association of vertnin and reveals additional QTL. *BMC Genetics*, 16, 129. <https://doi.org/10.1186/s12863-015-0286-9>
- Rousseau S. et al. (2013). A Genome-wide association study points out the causal implication of SOX9 in the sex-reversal phenotype in XX pigs. *PLoS ONE*, 8(11), e79882. <https://doi.org/10.1371/journal.pone.0079882>
- Schumacher T. et al. (2021). Congenital splay leg syndrome in piglets - current knowledge and a new approach to etiology. *Frontiers in Veterinary Science*, 8, 609883. <https://doi.org/10.3389/fvets.2021.609883>
- Sevillano C.A. et al. (2015). Genome-wide association study using deregressed breeding values for cryptorchidism and scrotal/inguinal hernia in two pig lines. *Genetics Selection Evolution*, 47, 18. <https://doi.org/10.1186/s12711-015-0096-6>
- Stenberg H. et al. (2020). A review of congenital tremor type A-II in piglets. *Animal Health Research Reviews*, 21, 84–88. <https://doi.org/10.1017/S146625232000002X>
- Szczerbal I. et al. (2019). Elevated incidence of freemartinism in pigs detected by droplet digital PCR and cytogenetic techniques. *Livestock Science*, 219, 52–56. <https://doi.org/10.1016/j.livsci.2018.11.009>
- Tomboc M. et al. (2000). Insulin-like 3/Relaxin-like factor gene mutations are associated with cryptorchidism. *The Journal of Clinical Endocrinology & Metabolism*, 85(11), 4013–4018. <https://doi.org/10.1210/jcem.85.11.6935>
- Wiedemann S. et al. (2005). Genome-wide scan for anal atresia in swine identifies linkage and association with a chromosome region on *Sus scrofa* chromosome 1. *Genetics*, 171(3), 1207–1217. <https://doi.org/10.1534/genetics.104.032805>
- Wu T. et al. (2018). Transcriptome analysis reveals candidate genes involved in splay leg syndrome in piglets. *Journal of Applied Genetics*, 59, 475–83. <https://doi.org/10.1007/s13353-018-0454-5>
- Xu S. et al. (2018). Polymorphisms of HOMER1 gene are associated with piglet splay leg syndrome and one significant SNP can affect its intronic promoter activity *in vitro*. *BMC Genetics*, 19, 110. <https://doi.org/10.1186/s12863-018-0701-0>
- Xu W. et al. (2019). Rediscover and refine QTLs for pig scrotal hernia by increasing a specially designed F3 population and using whole-genome sequence imputation technology. *Frontiers in Genetics*, 10, 890. <https://doi.org/10.3389/fgene.2019.00890>
- Žáková E. et al. (2020). System of collection and storage of health data in the performance testing of pigs. Certified methodology. Prague: Institute of Animal Science, 17 p. <http://vuzv.cz/en/publications/system-sberu-a-uchovani-zdravotnich-dat-v-kontrolu-uzitkovosti-prasat>. In Czech.